arXiv:hep-ph/0210067v1 4 Oct 2002

T violation in $\Lambda_b \to \Lambda \ell^+ \ell^-$ decays

Chuan-Hung Chen[a], C. Q. Geng[b] * and J. N. Ng [c]

[a]Institute of Physics, Academia Sinica
Taipei, Taiwan 115, Republic of China

[b]Department of Physics, National Tsing Hua University
Hsinchu, Taiwan, Republic of China

[c]Theory Group, TRIUMF, 4004 Wesbrrok Mall
Vancouver, B.C. V6T 2A3, Canada

We examine the T-odd transverse lepton and Λ polarizations in the baryonic decays of $\Lambda_b \to \Lambda l^+ l^-$ $(l = e\,, \mu)$ to study the T violating effects. We show that the lepton polarizations are suppressed but the Λ ones can be as large as 50% in CP violating theories beyond the standard model such as the SUSY models, which can be tested in various future hadron colliders.

Recently, time-reversal violation (TV) has been measured experimentally in the K^0 system [1], and thus complements the information on CP violation (CPV) that has been steadily accumulating for the past thirty seven years. The data is in accordance with the CPT theorem which is fundamental to local quantum field theories with Lorentz invariance and the usual spin-statistics connection. However, the origin of the violation remains unclear. In the standard model, CPV or TV arises from a unique physical phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [2]. This paradigm also predicts CPV effect in the b-quark system. To test the accuracy of this paradigm and to search for other sources of CPV one needs to look for new processes. Indeed this is an important quest of the B-factories. In addition we deem it particularly interesting if the time reversal symmetry violation can be directly detected in the b-system rather than inferring it as a consequence of CPT invariance. In this talk, we will present our recent studies on T violation in the baryonic decays of $\Lambda_b \to \Lambda l^+ l^-$ $(l = e, \mu)$ [3,4,5,6].

It is known that for a general three-body decay of a baryon the triple spin-momentum correlations, such as $\vec{s}_k \cdot (\vec{p}_i \times \vec{p}_j)$, are T-odd observables, where $\vec{s}_k$ and $\vec{p}_{i,j}$ are the spin and momentum vectors of the final particles, respectively. There are a number of different sources that might give rise to these T-odd observables. The most important ones being the weak CPV such as the CKM phase of the SM. However, final state interactions such as QCD for non-leptonic decays or the electromagnetic (EM) interaction among the final state particles can also make contributions. These are usually less interesting and they could even hide the signals from the weak CPV. We note that the T-odd triple correlations do not need non-zero strong phases unlike some of CP violating observables, such as the rate asymmetry between a particle and its antiparticle. In $\Lambda_b \to \Lambda l^+ l^-$ decays, we can use either the polarization of the lepton (s_l) or the Λ baryon (s_Λ) to study the T-odd correlations. It is interesting to note that the decays can be used to analyze the helicity structure of the interaction, which is impossible in the B meson system.

We start with the effective Hamiltonian for $b \to s l^+ l^-$ by including the right-handed couplings in

---

*Talk presented by C.Q. Geng at the 5th International Conference on Hyperons, Charm and Beauty Hadrons the hadronic sector are given by

$$\mathcal{H}(b \to sl^+l^-) = \frac{G_F \alpha_{em}}{\sqrt{2}\pi} V_{tb} V_{ts}^* (H_{1\mu} L_V^\mu + H_{2\mu} L_A^\mu)$$

$$\begin{aligned} H_{1\mu} &= \bar{s}\gamma^\mu (C_9^L P_L + C_9^R P_R) b \\ &\quad - \frac{2m_b}{q^2} \bar{s} i \sigma_{\mu\nu} q^\nu (C_7^L P_R + C_7^R P_L) b \,, \\ H_{2\mu} &= \bar{s}\gamma^\mu (C_{10}^L P_L + C_{10}^R P_R) b \,, \\ L_V^\mu &= \bar{l}\gamma^\mu l \,, \quad L_A^\mu \;=\; \bar{l}\gamma^\mu \gamma_5 l \,, \end{aligned} \tag{1}$$

where C_i^L and C_i^R $(i = 7, 9, 10)$ denote the effective Wilson coefficients of left- and right-handed couplings, respectively. In the standard model,

$$\begin{aligned} C_9^L &= C_9^{eff} \,, \qquad C_9^R = 0 \,, \\ C_{10}^L &= C_{10} \,, \qquad C_{10}^R = 0 \,, \\ C_7^L &= C_7^{eff} \,, \qquad C_7^R = \frac{m_s}{m_b} C_7^{eff} \,, \end{aligned} \tag{2}$$

where C_9^{eff}, C_{10}, and C_7^{eff} are the standard Wilson coefficients [7].

To study the exclusive decays of $\Lambda_b \to \Lambda l^+ l^-$, one needs to know the form factors in the transition of $\Lambda_b(p_{\Lambda_b}) \to \Lambda(p_\Lambda)$, parametrized generally as follows:

$$\begin{aligned} & \langle \Lambda | \, \bar{s} \; \Gamma_\mu \; b \, | \Lambda_b \rangle = f_1^{(T)} \bar{u}_\Lambda \gamma_\mu u_{\Lambda_b} \\ + \;& f_2^{(T)} \bar{u}_\Lambda i \sigma_{\mu\nu} \; q^\nu u_{\Lambda_b} + f_3^{(T)} q_\mu \bar{u}_\Lambda u_{\Lambda_b} \,, \\ & \langle \Lambda | \, \bar{s} \; \Gamma_\mu \gamma_5 \; b \, | \Lambda_b \rangle = g_1^{(T)} \bar{u}_\Lambda \gamma_\mu \gamma_5 u_{\Lambda_b} \\ + \;& g_2^{(T)} \bar{u}_\Lambda i \sigma_{\mu\nu} \; q^\nu \gamma_5 u_{\Lambda_b} + g_3^{(T)} q_\mu \bar{u}_\Lambda \gamma_5 u_{\Lambda_b} \,, \end{aligned} \tag{3}$$

where $\Gamma_\mu = \gamma_\mu$ $(i\sigma_{\mu\nu})$, and $f_i^{(T)}$ and $g_i^{(T)}$ are the form factors of vector (tensor) and axial-vector (axial-tensor) currents, respectively. In the heavy quark effective theory (HQET) [8], the form factors in Eq. (3) can be simplified by using

$$\begin{aligned} & \langle \Lambda(p_\Lambda) | \, \bar{s} \Gamma b \, | \Lambda_b(p_{\Lambda_b}) \rangle \\ = \;& \bar{u}_\Lambda \left(F_1(q^2) + \not{v} F_2(q^2) \right) \Gamma u_{\Lambda_b} \,, \end{aligned} \tag{4}$$

where Γ denotes the Dirac matrix, $v = p_{\Lambda_b}/M_{\Lambda_b}$ is the four-velocity of Λ_b, and $q = p_{\Lambda_b} - p_\Lambda$ is the momentum transfer, and the relations among the form factors can be found in Ref. [3]. Explicitly, under the HEQT, we have

$$f_1 \;=\; g_2 = f_2^T = g_2^T = F_1 + \sqrt{r} F_2 \,,$$

$$\begin{aligned} \rho \;&\equiv\; M_{\Lambda_b} \left(\frac{f_2 + g_2}{f_1 + g_1} \right) = \frac{M_{\Lambda_b}}{q^2} \left(\frac{f_1^T + g_1^T}{f_1 + g_1} \right) \\ &= \frac{F_2}{F_1 + \sqrt{r} F_2} \,. \end{aligned} \tag{5}$$

From experiments [9] and theories [8,10], one finds that $|F_1/F_2| > 1$ and $|\rho| < 1$.

To explore the T violating effects using the lepton or Λ spin polarization, we write i $(i = l$ or $\Lambda)$ four-spin vector in terms of a unit vector, $\hat{\xi}^i$, along the spin in its rest frame, as

$$s_{i0} = \frac{\vec{p}_i \cdot \hat{\xi}_i}{M_i}, \qquad \vec{s}_i \;=\; \hat{\xi}_i + \frac{s_{i0}}{E_i + M_i} \vec{p}_i, \tag{6}$$

and choose the unit vectors along the longitudinal, normal, transverse components of the i polarization, to be

$$\begin{aligned} \hat{e}_L^i \;&=\; \frac{\vec{p}_i}{|\vec{p}_i|}, \; \hat{e}_N^i \;=\; \frac{\vec{p}_i \times (\vec{p}_{l^-} \times \vec{p}_\Lambda)}{|\vec{p}_i \times (\vec{p}_{l^-} \times \vec{p}_\Lambda)|}, \\ \hat{e}_T^i \;&=\; \frac{\vec{p}_{l^-} \times \vec{p}_\Lambda}{|\vec{p}_{l^-} \times \vec{p}_\Lambda|} \,, \end{aligned} \tag{7}$$

respectively. Hence, the differential decay rates with polarized i is given by

$$d\Gamma^i(t) \;=\; \frac{1}{2} d\Gamma^0(t) \left[1 + \vec{P}^i \cdot \hat{\xi}^i \right] \,, \tag{8}$$

where $t = E_\Lambda / M_{\Lambda_b}$, $\vec{P}^i$ is the $i = l$ or Λ polarization vector, defined by

$$\vec{P}^i = P_L^i \hat{e}_L^i + P_N^i \hat{e}_N^i + P_T^i \hat{e}_T^i \,, \tag{9}$$

and $d\Gamma^0(t)$ is given in Ref. [6]. With the T odd transverse i polarization defined by

$$P_T^i \;=\; \frac{d\Gamma^i \left(\hat{\xi}_i \cdot \hat{e}_T^i = 1 \right) - d\Gamma^i \left(\hat{\xi}_i \cdot \hat{e}_T^i = -1 \right)}{d\Gamma^i \left(\hat{\xi}_i \cdot \hat{e}_T^i = 1 \right) + d\Gamma^i \left(\hat{\xi}_i \cdot \hat{e}_T^i = -1 \right)},$$

we obtain

$$\begin{aligned} P_T^l \;&\propto\; \frac{m_l}{M_{\Lambda_b}} Im \left(C_9^L C_{10}^{L*} + C_9^R C_{10}^{R*} \right) \,, \\ & \quad\; \frac{m_l}{M_{\Lambda_b}} Im \left(C_7^L C_{10}^{L*} + C_7^R C_{10}^{R*} \right) \end{aligned} \tag{10}$$

$$\begin{aligned} P_T^\Lambda \;&\propto\; Im \left(C_9^R C_{10}^{L*} - C_9^L C_{10}^{R*} \right) \,, \\ & \quad\; Im \left(C_7^L C_{10}^{R*} - C_7^R C_{10}^{L*} \right) \end{aligned} \tag{11}$$

It is easy to see that from Eq. (10) P_T^l is suppressed since it is always associated with the small lepton mass and to have a non-zero value of P_T^Λ, from Eq. (11), it is necessary to have conditions of (a) the existence of C_k^R and (b) a phase of $C_i^L C_j^R$ $(i \neq j)$. We remark that these conditions for P_T^Λ are clearly different from those of the T odd transverse lepton polarizations in both inclusive decay of $b \to s l^+ l^-$ [11] and exclusive ones, such as $B \to K^{(*)} l^+ l^-$ [12] and $\Lambda_b \to \Lambda l^+ l^-$ [4,5]. Such property distinguishes the T odd transverse Λ polarizations from various other T odd observables.

In the standard model, since there are no C_9^R and C_{10}^R as seen from Eq. (2), we have

$$\begin{aligned} P_T^l &\propto \frac{m_l}{m_b} Im\left(C_7^{eff} C_{10}\right) \\ P_T^\Lambda &\propto \frac{m_s}{m_b} Im\left(C_7^{eff} C_{10}\right) \end{aligned} \tag{12}$$

which are all suppressed. We note that the contribution to P_T^i from the EM final state interaction are $< O(10^{-3})$. Moreover, the long-distance (LD) effects in the one-loop matrix elements of $O_{1,2}$ [7] and $\Lambda_b \to \Lambda J/\Psi$ with $J/\Psi \to l^+ l^-$ are absorbed to C_9^L. On the other hand, it is clear that a large value of P_T^Λ, according to Eq. (11), can be obtained if a theory contains C_9^R or C_{10}^R or a large C_7^R, with a non-zero phase. Many theories beyond the standard model could give rise to C_9^R or C_{10}^R; examples are the left-right symmetric and supersymmetric models.

To illustrate our result, we use SUSY models [13,14] both with and without R-parity as shown in Ref. [6]. In Figure 1, we give the differential branching ratios (BRs) of $\Lambda_b \to \Lambda \mu^+ \mu^-$ with respect to E_Λ with and without including resonant states of Ψ and Ψ', and we find that the integrated BRs for the latter are (2.10, 1.66, 4.41) $\times 10^{-6}$ for the standard and SUSY with and without R-parity models, respectively. In Figures 2 and 3, we show $P_T^\Lambda(\Lambda_b \to \Lambda \mu^+ \mu^-)$ as a function of $E_\Lambda / M_{\Lambda_b}$ for the two types of models, with and without the LD contributions, respectively. As seen from the figures, even though the derivations of BRs to the standard model result are insignificant, the transverse Λ polarization asymmetries can be over 50% in both SUSY models with and without R-parity. Similar results are also expected for the decay of $\Lambda_b \to \Lambda e^+ e^-$. We remark that measuring a large P_T^Λ in $\Lambda_b \to \Lambda l^+ l^-$ is a clean indication of T violation as well as new CP violation mechanism beyond the standard model. To measure $P_T^\Lambda(\Lambda_b \to \Lambda \mu^+ \mu^-) \sim 10\%$ at 3σ level, at least 4.5×10^7 Λ_b decays are required if we use $BR(\Lambda_b \to \Lambda \mu^+ \mu^-) \sim 2 \times 10^{-6}$. Clearly, the measurement could be done in the second generation of B-physics experiments, such as LHCb, ATLAS, and CMS at the LHC, and BTeV at the Tevatron, which produce $\sim 10^{12} b\bar{b}$ pairs per year [15]. This is certainly within reach of a super B factory under discussion now [16]. Finally, we note that similar results are expected for using polarized Λ_b instead of Λ.



Figure 1. Differential BRs of $\Lambda_b \to \Lambda \mu^+ \mu^-$ as a function of E_Λ with and without resonant shapes, where the solid, dash-dotted and dashed curves stand for the results of the standard and SUSY with and without R-parity models, respectively.

Acknowledgments This work was supported in part by the National Center for Theoretical Science, National Science Council of the Republic



Figure 2. Transverse Λ polarization in $\Lambda_b \to \Lambda\mu^+\mu^-$ as a function of E_Λ/M_{Λ_b} in the SUSY model with R-parity. The curves with and without resonant shapes represent including and no long-distance contributions, respectively.



Figure 3. Same as Figure 2 but for the SUSY model without R-parity.

of China under Contract Nos. NSC-90-2112-M-001-069 and NSC-90-2112-M-007-040, and National Science and Engineering Research Council of Canada.

REFERENCES

1. CPLEAR Collaboration, A. Angelopoulos et al., *Phys. Lett.* **B444**, 43 (1998).
2. N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
3. C. H. Chen and C. Q. Geng, Phys. Rev. D **63**, 114024 (2001) [arXiv:hep-ph/0101171].
4. C. H. Chen and C. Q. Geng, Phys. Lett. B **516**, 327 (2001) [arXiv:hep-ph/0101201].
5. C. H. Chen and C. Q. Geng, Phys. Rev. D **64**, 074001 (2001) [arXiv:hep-ph/0106193].
6. C. H. Chen, C. Q. Geng and J. N. Ng, Phys. Rev. D **65**, 091502 (2002).
7. G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Rev. Mod. Phys.* **68**, 1230 (1996).
8. T. Mannel, W. Roberts and Z. Ryzak *Nucl. Phys.* **B355**, 38 (1991); T. Mannel and S. Recksiegel *J. Phys.* **G24**, 979 (1998).
9. CLEO Collaboration, G. Crawford *et al Phys. Rev. Lett.* **75**, 624 (1995).
10. C.S. Huang and H.G. Yan, *Phys Rev.* **D59**, 114022 (1999).
11. F. Kruger and L.M. Sehgal, *Phys. Lett.* **B380**, 199 (1996).
12. C.Q. Geng and C.P. Kao, *Phys. Rev.* **D57**, 4479 (1998).
13. F. Gabbiani et. al., Nucl. Phys. **B477**, 321 (1996); E. Lunghi et. al., *ibid* **B568,** 120 (2000).
14. D. Guetta and E. Nardi, *Phys. Rev.* **D58**, 012001 (1998) and references therein.
15. N. Harnew, Proc. of Heavy Flavors 8, (Southampton. 1999).
16. See, *e.g.*, G. Eigen, arXiv:hep-ex/0112042.